

Lecture 13 - M3-nets

Tuesday, February 18, 2020 14:37

today: M³-net trick: M vs. L polytope

II) M³-net example (CRF score)

* graph $G = (V, E)$

set of cliques \mathcal{C}
MRF model for score

$$y_p, p \in V \quad y_c \triangleq (y_p)_{p \in C}$$

$$p(y|x; w) = \frac{1}{Z(x; w)} \prod_{C \in \mathcal{C}} N_C(y_C; x, w)$$

$$= \frac{1}{Z} \exp \left(\sum_{C \in \mathcal{C}} \underbrace{\log N_C(y_C; x, w)}_{S(x, y; w)} \right)$$

$$w^T \varphi_C(y_C; x)$$

Example: CR
chain graph on y

$$\text{inference: } \max_{\tilde{y} \in S} \sum_{C \in \mathcal{C}} w^T \varphi_C(y_C; x)$$


goal: let's rewrite as a LP

* fix $x \in \mathbb{R}^n$; let S be a vector of size $|S(x)|$ of scores

$$\max_{\tilde{y} \in S(x)} s(x, \tilde{y}; w) = \max_k s_k = \max_{\{e_k\}} \langle e_k, s \rangle$$

$$\begin{aligned} \text{trick} \Rightarrow & \left(\begin{array}{l} \text{tight} \\ \text{convex} \\ (\text{vs. relaxation}) \end{array} \right) \stackrel{\rightarrow}{=} \max_{\alpha \in \text{conv}(\{e_k\})} \langle \alpha, s \rangle \\ & \stackrel{\Delta}{=} \Delta \alpha \end{aligned}$$

$$e_k = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} \xrightarrow{k^{\text{th position}}}$$



Note: Any comb. opt. problem [trivially] can be written as a convex opt. (LP) [but with exp # of dimensions]

⊕ insight of M³-net paper \rightarrow use MRF structure

$$\text{to transform } \max_{\alpha \in \Delta \Omega} \langle \alpha, s \rangle = \max_{M \in L} \sum_{C \in C} \langle M_C, s_C \rangle$$

reasonable size
tractable

marginal polytope (M) formulation

$$\begin{aligned} \langle \alpha, s \rangle &= \sum_{\tilde{y} \in S} \alpha(\tilde{y}) \left(\sum_{c \in C} s_c(\tilde{y}_c) \right) = \sum_{c \in C} \sum_{\tilde{y} \in S} \alpha(\tilde{y}) s_c(\tilde{y}_c) \\ &= \sum_{c \in C} \sum_{y' \in S_c} s_c(\tilde{y}_c) \left[\sum_{\substack{y \in S \\ y'_c = y_c}} \alpha(y) \right] \\ &\quad \triangleq \mu_c(\tilde{y}_c) \quad \text{"clique marginal variable"} \end{aligned}$$

$$\boldsymbol{\mu} \triangleq (\mu_c)_{c \in C} \quad \mu_c \in \Delta_{|S_c|}$$

$$M = A_M \text{ or}$$

$$\max_{\alpha \in \Delta_{|S|}} \langle \alpha, s \rangle = \max_{M \in A_M} \sum_{c \in C} \sum_{\substack{y \in S \\ y'_c = y_c}} s_c(y_c) \mu_c(\tilde{y}_c)$$

$$M \triangleq A_M \Delta_{|S|} \quad M \triangleq \left\{ (\mu_c)_{c \in C} : \exists \alpha \in \Delta_{|S|} \text{ s.t. } \mu_c(\tilde{y}_c) = \sum_{\substack{y \in S \\ y'_c = y_c}} \alpha(y) \forall c \right\}$$

$$M \subseteq \bigcup_{c \in C} \Delta_{|S_c|}$$

also $M \subseteq L \triangleq \{ (\mu_c)_{c \in C} : \mu_c \in \Delta_{|S_c|} \text{ and } \mu_c \text{'s are "locally consistent"} \}$

"local consistency
polytope"

i.e. $\forall C, C' \in \mathcal{C}$

$$\forall S \subseteq C \cap C'$$


$$\forall y_S$$


$$\sum_{\substack{y'_i \in S \\ \text{s.t. } y'_i = y_S}} \mu_c(y'_i) = \sum_{\substack{y'_i \in S \\ \text{s.t. } y'_i = y_S}} \mu_{c'}(y'_i)$$

$$= "L(y_S)"$$

(*) L can be seen as a relaxation of M

"If G is not triangulated, then we can have $L \not\subseteq M$





\otimes Key property: If G is triangulated then $M = L \Leftrightarrow \otimes \otimes$


$$\text{Implication here } \max_{M \in M} \left\{ \sum_{C \in C} l_C, n_C \right\} = \max_{L \in L} \quad \quad \quad \quad$$

$\underbrace{\qquad\qquad\qquad}_{g(w, z)}$


[assuming $l(y, y') = \sum l_C(y, y'_C)$]

aside: triangulated graph review:

Δ -graph \rightarrow iff it has no cycle of size 4 or more which are not chordal



\otimes you can triangulate a graph by running variable elimination




\otimes If G is triangulated \Rightarrow can construct a junction tree


\hookrightarrow Clique tree : tree with nodes being the with the running intersection property maximal cliques of G

\hookrightarrow if $v \in C_1 \cap C_2$ then v belongs to all cliques along the unique path from C_1 to C_2 in J.T.


Ex. of J.T. for G



\otimes to build a J.T., first build weighted complete graph on cliques where weight on edge $\{i,j\}$ is $|C_i \cap C_j|$



use maximum weight spanning tree to get J.T.





$\text{no } 3 \Rightarrow \text{no good! no q.J.T.}$

$M = L$ for Δ -graphs

• $M \subseteq L$

$$\bullet L \subseteq M : \alpha(y) = \frac{\prod_{c \in S} \mu(y_c)}{\prod_{S \in T} \mu_S(y_S)}$$

↓
Separator sets
from J.T.

Suppose $\mu \in L$;


We want $\exists \alpha$ s.t. μ is mang. of α

i.e. $\mu = \alpha \circ \varphi$

(this is gen.
that

for the free UGM

$$p(x) = \prod_{\substack{i,j \in E \\ p(x_i, x_j)}} \frac{p(x_i, x_j)}{p(x_i)p(x_j)} \prod_{i \in V} p(x_i)$$



Let's say want to show
that $M_B(y_B) = \sum_{y' : y'_B = y_B} \alpha(y')$

$$\sum_{y' : y'_B = y_B} \left(\frac{\prod_{c \in S} \mu_c(y'_c)}{\prod_{S \in T} \mu_S(y'_S)} \right)$$

There is no y_C here
since $y_C \notin S_{A,B}$
crucially using
non-inter. property

$$\text{now, } \sum_{y_A} M_A(y_A) = M_{\{2,3\}}(y_{\{2,3\}}) \quad [\text{by local consistency}]$$

$$= M_{S_{A,B}}(y_{S_{A,B}}) \quad |$$

$$y_i = \mu_{S_{A,B}}(y_{S_{A,B}})$$

keep plucking the leaves
until you are left with $\mu_B(y_B)$

Q) Why do we care to have shown $M=L$ for Δ -graph?

→ because describing L is simple i.e. $\text{Poly}(\max|S|) \# \text{of constraints}$

Vs. M which needs exponential # II in worst case

To get back to M^2_{net} :

$$\max_{\tilde{y} \in S} l(\tilde{y}) = \max_{\alpha \in \Delta^{|\mathcal{X}|}} \langle \alpha, S \rangle = \max_{M \in \mathcal{M}} \sum_{C \in \mathcal{C}} \langle M_C, S_C \rangle = \max_{M \in \mathcal{M}} \sum_{C \in \mathcal{C}} \langle M_C, S_C \rangle$$

when $M=L$ (i.e. Δ -graph)



Note:
could use
junction tree dly
[max form]
(generalization of
trivbi)

to solve this LP
much more efficiently
(than interior point method)

$$\text{all the vertices have integer components (i.e. } \{0,1\} \text{)} \\ \text{i.e. } \mu^* = \arg \max_{\mu \in \mathcal{L}} \sum_{C \in \mathcal{C}} \langle \mu_C, S_C \rangle \\ \text{will have } \mu_C^*(\cdot) = \delta_{y_C^*}(\cdot)$$

Overall, suppose that $l_i(y)$ decomposes also on G

$$\max_{\tilde{y} \in S_i} l_i(\tilde{y}) - w^T \Phi_i(\tilde{y}) = \max_{\mu \in \mathcal{L}_i} a_i + (c_i + F_i^T w)^T \mu - w^T F_i \mu$$

(integer) marginal
ground truth y^*

Last class: A) dualize this LP and plug it in structured SVM obj.
to get a small "complicated" LP

use Nesterov, CPLEX, etc. to solve to 1e-6 accuracy use IPM e.g.

B) saddle pt. formulation $\min_w \max_{\mu \in \mathcal{L}_i} g(w, \mu)$

extragradient → would require projection on \mathcal{L}_i
which is more expensive than LP

c) later: saddle pt. FW alg which only requires LMOs
"linear min. oracles"
on L^1_S
instead of projection

next class: Lagrangian dual of SVM struct