


Lecture 14 - dual SVM

Thursday, February 27, 2020 13:42

- today:
- IPM
 - SVM dual



interior point method:

$$\min f(x)$$


$$\text{s.t. } Ax = b \\ x \geq 0$$

barrier method

$$\min f(x) + t(\frac{1}{t} \log x) \stackrel{m}{=} g_t(x)$$

$$\text{s.t. } Ax = b \\ \frac{1}{t} \log x \geq 0$$

$$\text{dom}(g_t(x)) = \{x : x \geq 0\}$$



$$x_t^* = \underset{\text{s.t. } Ax = b}{\operatorname{arg\,min}} g_t(x)$$

Interior pt. method: run Newton's method with warm start on $g_t(x)$ and decrease t with geometric schedule "self-concordant analysis"

Convergence: Newton's method take $\log(\log(\frac{1}{\epsilon})) + \text{const.} (\|f(x_0) - f^*\|)$ iterations to reach ϵ -accuracy

IPM: # of Newton iterations is $\log(\frac{1}{\epsilon})$. same constant overall convergence

↳ dimensionality is appearing; but not condition # of f (vs. gradient method)

Review of Lagrangian duality

primal problem

$$P^* \triangleq \inf_{\mathbf{x}} f(\mathbf{x}) \\ \text{s.t. } h_i(\mathbf{x}) \leq 0 \quad \forall i \in I$$

dual problem

$$d^* \triangleq \sup_{\alpha \geq 0} d(\alpha)$$

Lagrangian dual
sol.

Fenchel conjugate of f

$$f^*(y) \triangleq \sup_{\mathbf{x}} \langle y, \mathbf{x} \rangle - f(\mathbf{x})$$

$$P_{\mathbf{x}} \text{ s.t. } \nabla f(\mathbf{x}) = y$$

Fenchel duality:

$$P^* = \inf_{\mathbf{x}} f(\mathbf{x}) + g(A\mathbf{x}) \quad A: \mathbb{R}^d \rightarrow \mathbb{R}^P \\ \frac{\|w\|^P}{2} + \sum_{i=1}^n h_i(w) \\ d^* = \sup_y -f^*(A^T y) - g^*(y)$$

$$\max_w l(w, y) \\ \tilde{g} = -\nabla w, P_i(y)$$

Note: not unique?

middle of. perspective for dual duality

Lagrange multipliers

Note, not unique:

saddle pt. perspective for lag. duality

$$\inf_{\mathbf{x}} \underset{\mathbf{h}(\mathbf{x}) \leq 0}{f(\mathbf{x})} = \inf_{\mathbf{x}} \left[\sup_{\alpha \geq 0} f(\mathbf{x}) + \underbrace{\sum_i \alpha_i h_i(\mathbf{x})}_{L(\mathbf{x}, \alpha)} \right] \quad \text{Lagrange multiplier}$$

"Lagrangian fct"

$$\text{because } g(\mathbf{x}) \triangleq \sup_{\alpha \geq 0} \sum_i \alpha_i h_i(\mathbf{x}) = \begin{cases} +\infty & \text{if } \mathbf{x} \text{ is not feasible} \\ 0 & \text{otherwise} \end{cases}$$

G.W.
where $C \triangleq \{\mathbf{x} : h_i(\mathbf{x}) \leq 0 \forall i \in I\}$

$$p^* = \inf_{\mathbf{x}} \sup_{\alpha \geq 0} L(\mathbf{x}, \alpha)$$

$$\geq \sup_{\alpha \geq 0} \inf_{\mathbf{x}} L(\mathbf{x}, \alpha) \triangleq d^*$$

"weak duality"
Lagrangian dual fct.

Note: $L(\mathbf{x})$ always concave

even if $f(\mathbf{x})$ is not convex

$\inf_{\mathbf{x} \in C} \geq \sup_{\alpha \geq 0} \inf_{\mathbf{x}}$?

Strong duality $p^* = d^*$

$$\sup_{\alpha} L(\mathbf{x}, \alpha) \geq \inf_{\mathbf{x}} f(\mathbf{x})$$

KKT conditions: \rightarrow necessary conditions for $\mathbf{x}^* \in \mathcal{X}$ to be primal & dual optimum resp. when $f \& h_i$ are diff. fcts.



- 14h44
- KKT conditions
- \mathbf{x}^* primal feasible (*i.e.* $h_i(\mathbf{x}^*) \leq 0 \forall i \in I$)
 - α^* dual feasible ($\alpha \geq 0$)
 - $\nabla f(\mathbf{x}^*) + \sum_i \alpha_i^* \nabla h_i(\mathbf{x}^*) = 0$ [*i.e.* (\mathbf{x}^*, α^*) stationary p. of $L(\mathbf{x}, \alpha)$ with respect to \mathbf{x}]
 - $\alpha_i^* h_i(\mathbf{x}^*) = 0 \forall i$ complementary slackness

* If primal problem is convex
(*i.e.* $f \& h_i$'s are convex) and "Slater's condition" \Rightarrow strong duality *i.e.* $p^* = d^*$

$\exists \mathbf{x}$ rel. int. ($\text{dom}(f)$)
 $\exists i : h_i(\mathbf{x}) < 0$
when h_i is not affine

and KKT conditions are sufficient

(thus suppose get α^* which solves dual)
Then get \mathbf{x}^* through KKT
 $\Rightarrow \mathbf{x}^*$ sole primal

↳ counter-example for strong duality (see Boyd's book ex. 5.21)

$$\min f(\mathbf{x}, y) \quad \text{where } f(\mathbf{x}, y) = \begin{cases} -x/y & y > 0 \\ +\infty & \text{o.w.} \end{cases}$$

$\text{s.t. } \frac{x^2}{y} \leq 0$

1. on $x < 0$, $y < 0$

(Violates Slater's cond.)
and no strong duality

Other comments:

- $f(\alpha) \geq d(\alpha)$ $\forall \alpha \in \text{or feasible}$
- $d(\alpha)$ is always concave in α
- $d(\alpha)$ is obtained through uncons.-opt.
- No unique "Lagrangian dual problem" \rightarrow depends how you write primal problem
e.g. $h_i(z) \leq 0$ vs. $a_i h_i(z) \leq 0 \quad \forall a > 0$

SVM strict dual problem

Constrain primal: $\min_{w, \xi} \frac{\lambda \|w\|^2}{2} + \frac{1}{n} \sum_i \xi_i$
 s.t. $\frac{1}{n} \sum_i \xi_i \geq \sum_i H_i(\tilde{y}; w) \quad \forall \tilde{y} \in \mathbb{R}$
 $\downarrow l_i(\tilde{y}) - w^T \gamma_i(\tilde{y})$

$$L(w, \xi; \alpha) = \frac{\lambda \|w\|^2}{2} + \left(\sum_i \xi_i + \frac{1}{n} \sum_i \alpha_i (\tilde{y}_i) \underbrace{[l_i(\tilde{y}) - w^T \gamma_i(\tilde{y}) - \xi_i]}_{h_i(\tilde{y}; w, \xi)} \right)$$

$$\frac{\partial L}{\partial \xi_i} = 0 \Rightarrow \boxed{\sum_i \alpha_i l_i(\tilde{y}) = 1} \quad \text{i.e. } \alpha_i \in \Delta_{\{x_i\}}$$

$$\nabla_w L = 0 \Rightarrow \lambda w - \frac{1}{n} \sum_i \alpha_i l_i(\tilde{y}) \gamma_i(\tilde{y}) = 0$$

$$\boxed{w(\alpha) \triangleq \frac{1}{\lambda n} \sum_{i=1}^n \left(\sum_{\tilde{y} \in \{x_i\}} \alpha_i(\tilde{y}) \gamma_i(\tilde{y}) \right)} \quad \text{KKT condition}$$

$$d(\alpha) = \frac{\lambda}{2} \|w(\alpha)\|^2 + \underbrace{\left(\sum_i \alpha_i l_i(\tilde{y}) \right)}_{\langle b, \alpha \rangle} \left[l_i(\tilde{y}) - w(\alpha)^T \gamma_i(\tilde{y}) \right]$$

$$\begin{aligned} & \langle b, \alpha \rangle \\ & \text{rotation } \alpha \in \Delta_{\{x_1\}} \times \Delta_{\{x_2\}} \dots \times \Delta_{\{x_n\}} \\ & b = (b_i)_{i=1}^n \quad b_i(\tilde{y}) \triangleq \sum_i \alpha_i(\tilde{y}) \end{aligned}$$

$$d(\alpha) = \langle b, \alpha \rangle + \frac{\lambda}{2} \|w(\alpha)\|^2 - w(\alpha)^T \underbrace{\left(\sum_i \alpha_i l_i(\tilde{y}) \gamma_i(\tilde{y}) \right)}_{\Delta w(\alpha)}$$

$$d(\alpha) = \langle b, \alpha \rangle - \frac{\lambda}{2} \|w(\alpha)\|^2$$

$$\dim(\alpha) = m$$

$$b \in \mathbb{R}^m$$

$$A \in \mathbb{R}^{d \times m}$$

$$||w|| = \sqrt{b^T b - \frac{1}{2} ||w||^2}$$

dual problem:

$$\begin{aligned} & \text{max}_{\alpha_i \in A_{\tilde{y}, i}} -\frac{1}{2} \|w(\alpha)\|^2 + \langle b, \alpha \rangle \\ & \quad 1 \leq i \leq n \\ & \quad (\text{exponentially many dual variables}) \end{aligned}$$

DEIR

$$A \in \mathbb{R}^{d \times m}$$

$$w(\alpha) = A\alpha$$

$$A \triangleq \left[\cdots \begin{matrix} \alpha_1(\tilde{y}) \\ \vdots \\ \alpha_n(\tilde{y}) \end{matrix} \cdots \right]$$

i.e. column

$$\text{Complementary Slackness: } \alpha_i^*(\tilde{y}) > 0 \Rightarrow h_i(\tilde{y}; w^*) = 0$$

$$\text{i.e. } \xi_i^* = b_i(\tilde{y}) - w^T \gamma_i(\tilde{y}) = H_i(\tilde{y}; w^*)$$

$$\text{by primal feas., } \xi_i^* \geq H_i(\tilde{y}; w^*) \quad \forall \tilde{y} \in \mathcal{Y}$$

$$\text{thus } \xi_i^* = \max_{\tilde{y} \in \mathcal{Y}_i} H_i(\tilde{y}; w^*) = H_i(w^*)$$

Support vectors are when $\alpha_i^*(\tilde{y}) > 0$ [and $\tilde{y} \neq y^{(i)}$ because $\gamma_i(y^{(i)}) = 0$]

\circlearrowleft sparse solution α^*

[necessary condition $\tilde{y} \in \arg \max_{\tilde{y} \in \mathcal{Y}_i} H_i(\tilde{y}; w^*)$]