

Lecture 16 - FW

Thursday, March 12, 2020 13:35

- today:
 - away-step FW
 - FW convergence & properties

away-step FW

comment: $x_t = \sum_u p_u^t s_u$

$$x_{t+1} = (1-\gamma_t) x_t + \gamma_t s_t$$

$$= \sum_u p_u^{t+1} (1-\gamma_t) s_u + \gamma_t s_t$$

$p_u^{t+1} \rightarrow$ previous coordinates shrunk by $(1-\gamma_t)$

* FW step moves mass uniformly away from active set $\{s_u\}_{u=1}^t$ to FW corner s_t

→ unless step size $\gamma_t = 1$, FW never removes a corner from its expansion

⇒ zig-zag phenomenon close to boundary on polytopes




(this is why $\mathcal{O}(1/t)$ rate even if f is μ -strongly convex)

$$\left\langle -\nabla f(x_t), \frac{d_t}{\|\nabla f(x_t)\|} \right\rangle > \frac{\alpha_t}{\|d_t\|} \rightarrow 0 \text{ as } t \rightarrow \infty$$

⇒ slow rate

for strongly
convex?

$\mathcal{O}(\text{apl-pt})$
rate



* but FW has no problem — on "strongly convex sets"

↳ sublevel set of a strongly convex fct.

when sol'n
is in the relative
interior of M



)(


away-step FW fix: (solves zig-zagging problem)

in addition to compute

$$s_t = \operatorname{argmin}_{s \in M} \langle \nabla f(x_t), s \rangle$$

also

$$\text{compute } v_t \triangleq \operatorname{argmax}_{s \in \text{activeset}(x_t)} \langle \nabla f(x_t), s \rangle$$



also

$$\text{compute } v_t \triangleq \underset{\text{sc defined at } x_t}{\operatorname{argmax}} \langle \nabla f(x_t), s \rangle$$

$$d_{FW} \triangleq s_t - x_t$$

$$d_A \triangleq x_t - v_t$$



(aside: Wolfe's original alg. used what M here
⇒ non-convergent alg?)

* AFW picks the direction with best inner-product

$$\text{i.e. pick } d_A \text{ if } \underbrace{\langle d_A, -\nabla f(x_t) \rangle}_{\text{"g*"}} > \underbrace{\langle d_{FW}, -\nabla f(x_t) \rangle}_{\text{g*}}$$

or pick d_{FW}

d_A can be zero when $\nabla f(x_t)$ is \perp to active set

* if use d_A , let $\gamma_t = \underset{\gamma \in [0, \gamma_{\max}]}{\operatorname{argmin}} f(x_t + \gamma d_A)$

$$\gamma \in [0, \gamma_{\max}]$$

where γ_{\max} depends on A^t coefficients

$$\text{i.e. suppose } x_t = \sum_u \alpha_u s_u$$

$$x_{t+1} = x_t + \gamma_t (x_t - v_t)$$

$$= \sum_u ((1+\gamma_t) \alpha_u s_u - \gamma_t v_t)$$

let α be coefficient of v_t

$$(1+\gamma_{\max})\alpha - \gamma_{\max} = 0$$

$$\Rightarrow \boxed{\gamma_{\max} = \frac{\alpha}{1-\alpha}}$$

* when $\gamma_t = \gamma_{\max}$

"drop step" → remove v_t from expansion

⊕ when run AFW:

either you maintain some expansion of $x_t = \sum_u p_u s_u$

or

you have a feasibility oracle + away-step oracle


[see NIPS 2016 paper

by Meshi & Ganesh]

assumption to prove convergence

↳ they assume $\text{corners}(M) \subseteq \{x_0, \dots, x_n\}$





↳ they assume $\text{corners}(M) \subseteq \mathbb{S}_{U_1}$
 and M is described as $Ax = b$
 $x \geq 0$
 ↳ assumption to run alg.

AFW has linear convergence rate on polytopes when f is strongly convex

14h35

* combine d_{FW} & $d_A \rightarrow \text{pairwise FW}$

$$d_{\text{FW}} + d_A = s_t - x_t + x_t - v_t = \underbrace{s_t - v_t}_{d_{\text{PFW}}} \\ \underbrace{\langle -\nabla f(x_t), d_{\text{FW}} \rangle}_{g_{\text{FW}}} = g_{\text{FW}} + g_A$$

during AFW: $g_t = \langle -\nabla f(x_t), d_t \rangle = \max \{ g_{\text{FW}}, g_A \}$

$$g_t \geq \frac{g_{\text{FW}}}{2}$$

⊗ note: if $M = \text{conv}(\mathcal{A})$ where \mathcal{A} is some finite set (called "atoms")

$$\text{LMO}(r) : \min_{s \in M = \text{conv}(\mathcal{A})} \langle s, r \rangle = \min_{a \in \mathcal{A}} \langle a, r \rangle$$


↳ lot of application in ML



e.g. $\mathcal{A} \rightarrow$ integer flows
 $\text{conv}(\mathcal{A}) \rightarrow$ flow polytope LMO \rightarrow min cost network flow

$\mathcal{A} \rightarrow$ clique assignment in graph $(S_C)_{C \in \mathcal{F}}$ LMO \rightarrow max product alg.
 $\text{conv}(\mathcal{A}) \rightarrow$ marginal polytope or
 Graph-cut alg.

for submodular potentials
 (see later)



more properties of FW

6) convergence for non-convex f :

aside: necessary first order condition for constrained opt.

$$\min_{x \in M} f(x) \quad x^* \text{ is a local min}$$

$$\Rightarrow \langle \nabla f(x^*), S - x^* \rangle \geq 0 \quad \forall S \subset M$$

'stationpt.' for const.
 opt. problem



$\min_{x \in M} f(x)$ x^* is a local min \leftarrow minim pr. per const.
 $\Leftrightarrow \langle \nabla f(x^*), s - x^* \rangle \geq 0 \quad \forall s \in M$ opt. problem
 $\Leftrightarrow \min_{s \in M} \langle \nabla f(x^*), s - x^* \rangle \geq 0$
 $\Leftrightarrow \max_{s \in M} \langle -\nabla f(x^*), s - x^* \rangle \leq 0$ (If f and M are convex,
this is a sufficient cond.)
Fw-gap(x^*)
 \hookrightarrow quantify the "nonstationarity"

see L-J. 2016 and ir
 $\min_{s \in M} \text{gap}(x_s) \leq O(\frac{1}{\sqrt{t}})$ for FW with one search
 "non-convex" FW
 \int is L-smooth and lowerbound
 M is bounded & convex
but f is not nec. conv

4) affine covariance of FW

let \tilde{M} be a new domain s.t. $\tilde{M} \xrightarrow[A]{\text{union tract.}} M$ i.e. $M = A\tilde{M}$

define $\tilde{f}(\tilde{x}) \triangleq f(A\tilde{x})$

$$\min_{\tilde{x} \in \tilde{M}} \tilde{f}(\tilde{x}) = \min_{\tilde{x} \in \tilde{M}} f(A\tilde{x}) = \min_{\substack{x \in M \\ \tilde{x} \in \tilde{M}}} f(x)$$

affine covariance of FW:

if run FW on $\tilde{f} \notin \tilde{M}$ to get \tilde{x}_t it starts

then $x_t \triangleq A\tilde{x}_t$ corresponds to running FW on $f \in M$

(modulo tie-breaking)

$$\begin{array}{ccc} x_t & \xrightarrow{A} & \tilde{x}_t \\ \text{FW} \downarrow & & \downarrow \text{FW} \\ \tilde{f} \in M & \xrightarrow{A} & f \in \tilde{M} \end{array}$$

why? \rightarrow inner product is affine invariant $\langle \nabla_{\tilde{x}} \tilde{f}(\tilde{x}) \rangle = \langle \nabla_x f(A\tilde{x}) \rangle = A^T \langle \nabla_x f(x) \rangle$

$$\tilde{x}_t = \arg \min_{s \in M} \langle \tilde{s}, \nabla_{\tilde{x}} \tilde{f}(\tilde{x}_t) \rangle \quad x_t \triangleq A\tilde{x}_t$$

$$\langle \tilde{s}, A^T \nabla_x f(x_t) \rangle$$

$$\langle \underbrace{\tilde{A}\tilde{s}}_{s \in M}, \nabla_x f(x_t) \rangle$$

$$x_t = \arg \min_{\substack{s \in M \\ \tilde{A}\tilde{s}}} \langle s, \nabla_x f(x_t) \rangle$$

$$\begin{aligned} \tilde{x}_t &= \underline{A\tilde{x}_t} && (\text{modulo tie breaking}) \\ \tilde{s}_t &= \underline{A\tilde{s}_t} \end{aligned}$$

$$\text{SEM} \quad \overbrace{\text{A} \tilde{\text{M}}}^{\text{S}}$$

$$\overbrace{\text{D}_t}^{\text{S}} = \text{A} \tilde{\text{S}}_t$$

\Rightarrow we want affine invariant analysis $\rightarrow C_S \leq L_{11} (\det)_{1,1} (M)^2$