

Lecture 21 - scribbles

Tuesday, November 20, 2018 14:34

today: continue MCMC
Gibbs Sampling & MH.

continue Markov chain review:

Def: irreducible Markov chain \Leftrightarrow there exists a positive prob. "path"
from every $i \xrightarrow{*} j$

$$\forall (i, j), \exists \text{ an integer } m_{ij} \text{ s.t. } (A^{m_{ij}})_{ij} > 0$$

(by Perron-Frobenius thm.) \Rightarrow irreducible M.C. has a unique stationary dist.

* in order to converge to it, we need aperiodicity as well

irreducible and aperiodic M.C. \Leftrightarrow \exists an integer m s.t. $A^m > 0$

aka. regular M.C. (i.e. $(A^m)_{ij} > 0 \forall ij$)

(finite state)

or ergodic M.C.

(*) [note: a sufficient condition for an irreduc. M.C.
to be aperiodic is $\exists i$ s.t. $A_{ii} > 0$]

$$A = \begin{pmatrix} 0 & \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & 0 & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} & 0 \end{pmatrix} = \frac{1}{2}(J\mathbf{I}^T - \mathbf{I})$$

12 1 constant 1 1 1

$$A^2 = \frac{1}{4} \begin{pmatrix} 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 \end{pmatrix} \quad / \quad R$$

$$= \frac{1}{4} (R - 2AA^T + I)$$

Thm: If a finite M.C. is ergodic (regular)

then \exists a unique stationary dist. π

and for any starting dist. π_0 , $\lim_{t \rightarrow \infty} A^t \pi_0 = \pi$

the speed of convergence is related to the mixing time τ of the chain

$$\tau \triangleq \frac{1}{1 - |\lambda_2(A)|}$$

λ_2 2nd biggest eigenvalue of A

$$\|A^t \pi_0 - \pi\|_1 \leq C \exp(-t/\tau)$$

* intuition (from linear algebra) [informal argument]

basis
↓

simpler case, suppose A is diagonalizable with orthogonal matrix U
is symmetric

$$A = U \Sigma U^T \text{ with } \Sigma = \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \ddots & & \\ \vdots & & \ddots & 0 \\ 0 & \dots & 0 & \lambda_K \end{pmatrix}$$

(by Frobenius thm.)

$U \rightarrow$ basis of eigenvectors

$U = \{u_1, \dots, u_k\}$ take $u_i = \pi / \|u_i\|_2$ can show that $\lambda_1 = 1 > |\lambda_2| \geq \dots \geq |\lambda_k|$

Let α_o be coordinate in basis U st. $U^T U = I = U U^T$

$$\begin{aligned} \pi_o &= U \alpha_o \quad (\text{i.e. } \alpha_o = U^T \pi_o \quad (\alpha_o)_i = \frac{\pi^T \pi_o}{\|u_i\|_2}) \\ A^T \pi_o &= (U \cancel{U^T})^T \cancel{(U \cancel{U^T})} \dots (U \cancel{U^T})^T \cancel{U} \alpha_o \\ &= U \sum_{\alpha_o}^t \end{aligned}$$

$$A^T \pi_o = \left[(\alpha_o)_1 \overset{u_1}{\downarrow} \pi + (\alpha_o)_2 \overset{u_2}{\downarrow} \pi + \dots + (\alpha_o)_k \overset{u_k}{\downarrow} \pi \right] \overset{\rightarrow 0}{\underset{\lambda_2}{\downarrow}} \lambda_k^t$$

because $I^T A^T \pi_o = I \forall t \Rightarrow (\alpha_o)_1 = 1$ [not fully correct?]

$$\|A^T \pi_o - \pi\|_1 \leq C |\lambda_2|^t$$

$$|\lambda_2| = \underset{\text{first e-gap}}{\downarrow} -\varepsilon_1 \quad \varepsilon_1 \triangleq 1 - |\lambda_2|$$

$$|\lambda_2| \leq \exp(-\varepsilon_1) \quad 1-x \leq \exp(-x) \quad \forall x$$

$$|\lambda_2|^t \leq \exp(-t\varepsilon_1) \quad \Rightarrow \quad \gamma = \frac{1}{1 - |\lambda_2|} \quad //$$

* mixing time is often exponentially big

④ How do we design A st. $A^T \pi_0 \rightarrow \pi$?

one "easy way"

reversible M.C. \Leftrightarrow iff \exists dist. π s.t. $A_{ij}\pi_j = A_{ji}\pi_i \quad \forall (i,j)$

"detailed balance equation"

It means that $P\{X_t=i, X_{t-1}=j\} = P\{X_t=j, X_{t-1}=i\}$
(when $P\{X_{t-1}=i\} = \pi_i$)

sufficient condition (but not necessary)

for $A_{ii}=\pi_i$

proof: $(Aq)_i = \sum_j A_{ij}\pi_j \stackrel{\text{detailed balance}}{=} \sum_j A_{ji}\pi_i = \pi_i \left(\sum_j A_{ji} \right) //$

15h25

Metropolis-Hastings alg.

goal
construct a M.C. with stationary dist. $p(x)$ [constant]

(assume $p(x) > 0 \forall x$)

uses some proposal $q(x'|x)$

[i.e. if in state x , we sample $x' | x \sim q(x'|x)$.]

depends on $\frac{p(x')}{p(x)}$, no need to normalize

[i.e. if in state x , we sample $x' | x \sim q(x'|x)$]

accept new state x' with prob.

if reject \rightarrow stay in same state x

[this still a new sample]

vs. rejection sampling

where only "accepted states" are sampled

$$a(x'|x) \triangleq \min \left\{ 1, \frac{q(x|x') p(x)}{q(x'|x) p(x)} \right\}$$

acceptance ratio to satisfy detailed balance

MH alg.:

start at $x^{(0)}$

for $t=1, \dots$

• propose $x^{(t)} \sim q(x'|x^{(t-1)})$

• flip a biased coin with prob $a(x^{(t)} | x^{(t-1)})$ to be 1

• if accept (coin = 1):

let $x^{(t)} = x^{(t)}$

o.w.

$x^{(t)} = x^{(t-1)}$

end for

note: for symmetric $q(x'|x)$, always accept if $p(x') \geq p(x)$

[Metropolis alg.] \rightarrow like a noisy hill-climbing alg.

/target

[Metropolis alg.]

[verify as exercise that it satisfies detailed balance]

→ with $\pi = p$ ^{target}

* for convergence: if M.H. chain is ergodic, then we converge to correct unique stationary dist. p

sufficient conditions

for irreducibility $q(x'|x) > 0 \quad \forall x' \neq x \in X$

for aperiodicity $q(x|x) > 0$ for some $x \in X$

✳ aside: it is still ok to change proposal with time
(inhomogeneous M.C.) $q_t(x'|x)$

as long as choice of q_t does not depend on $x^{(t+1)}$


Then convergence theory above will go through


[i.e. detailed balance etc.
will give right stationary dist.]

slow mixing example:


Suppose p is a multivariate normal

$$\{ q(x'|x) = N(x'|x, \sigma^2 I)$$





* high prob. of rejection



here the ^{best.} mixing time is related to ratio $\frac{\sigma_{\max}}{\sigma_{\min}}$

good books: Casella & Berger

"Monte Carlo Statistical methods"


Gibbs sampling alg.:

↳ M.H. with a clever choice of proposal $q_t(x'|x)$

examples of applications:

→ U GM: $\tilde{p}(x) = \prod_c \psi_c(x_c)$

difficult conditional
in DGM



$$\tilde{p}(x) = p(x, \bar{x}_E) \delta(x_E, \bar{x}_E) \propto p(x | \bar{x}_E)$$

UGM:

cyclic Gibbs sampling alg.: nodes $i=1, \dots, n$

start at some $x^{(0)}$

for $i=1$

start at some x^0

for $t=1, \dots$

- pick $i = (t \bmod n) + 1$
- sample $x_i^{(t)} \sim p(x_i = \cdot | x_{\neq i}^{(t-1)})$
- set $x_j^{(t)} = x_j^{(t-1)}$ for $j \neq i$ time conditional as proposal

end

⊗ G.S. is M.H. with a time varying proposal

suppose we pick i at time t

for rest to be constant

$$\text{then proposal is } q_t(x' | x^{(t-1)}) = p(x'_i | x_{\neq i}^{(t-1)}) \delta(x'_{\neq i}, x_{\neq i}^{(t-1)})$$

acceptance ratio:

$$\alpha(x' | x^{(t-1)}) = \frac{p(x' | x^{(t-1)} | x')}{q_t(x' | x^{(t-1)} | x')} \frac{p(x')}{p(x^{(t-1)})}$$

$$= \frac{\cancel{q_t(x' | x^{(t-1)} | x')}}{\cancel{p(x'_i | x_{\neq i}^{(t-1)})}} \frac{\delta(x'_{\neq i}, x_{\neq i}^{(t-1)})}{\cancel{p(x_{\neq i}^{(t-1)})}} \frac{p(x'_{\neq i})}{\cancel{p(x'_i | x_{\neq i}^{(t-1)})}}$$

$$= 1 \quad \text{always accepts?}$$

Convergence of G.S.:

"

Convergence of G.S.:

Let A be Markov transition kernel of one full cycle of GS (i.e. n steps)

→ homog. M.C.

$$p(x) > 0 \forall x$$

→ A is irreducible & aperiodic because $A_{ij} > 0 \forall i, j$
 (since can get to any state with n steps)

$$\rightarrow \text{supposing } p(x_i | x_{\tau_i}) > 0 \quad \forall x_i \in \mathcal{X}$$


$$\rightarrow A^t \xrightarrow[t \rightarrow \infty]{\pi_0} P$$

⊕ also works for random scan (pick $i \sim \text{Unif}(\{n\})$ at each step)

example : G.-S. for Ising model

Ising model $x_i \in \{-1, 1\}$

UGM :



$$p(x) = \frac{1}{Z(n)} \exp \left(\sum_i m_i x_i + \sum_{i,j, j \in E} n_{ij} x_i x_j \right)$$

for G.S.,

want to compute $p(x_i | x_{\tau_i}) \propto p(x_i, x_{\tau_i})$

$$= \exp(m_i x_i + \sum_{j \in N(i)} n_{ij} x_i x_j + \text{rest})$$

- - 1 1 .. 1

\Rightarrow renormalize to get conditional

$$p(x_i=1 | x_{\bar{i}}) = \frac{\exp(\eta_i + \sum_{j \in N(i)} \eta_{ij} x_j)}{1 + \exp(-\eta_i - \sum_{j \in N(i)} \eta_{ij} x_j)} \cdot \frac{\exp(\alpha \delta)}{\exp(\alpha \delta)}$$
$$= \sigma(\eta_i + \sum_{j \in N(i)} \eta_{ij} x_j)$$