

# IFT 6760A - Lecture 2

## SVD and Orthogonality

**Scribe(s):** Yann BOUTEILLER, Parviz HAGGI

**Instructor:** Guillaume Rabusseau

### 1 Summary

In the previous lecture we reviewed the basics of linear algebra. In this lecture we will first look into the concept of Singular Value Decomposition of a matrix, often called SVD, followed by the concepts of orthogonality and orthogonal projections.

### 2 Singular Value Decomposition

**Theorem 1 (SVD).** Any matrix  $\mathbf{A} \in \mathbb{R}^{m \times n}$  can be written as:

$$\mathbf{A} = \mathbf{U}\mathbf{\Sigma}\mathbf{V}^T$$

where  $\mathbf{U} \in \mathbb{R}^{m \times m}$  and  $\mathbf{V} \in \mathbb{R}^{n \times n}$  are both orthogonal matrices (i.e.  $\mathbf{U}^T\mathbf{U} = \mathbf{I}_m$  and  $\mathbf{V}^T\mathbf{V} = \mathbf{I}_n$ ) and  $\mathbf{\Sigma} \in \mathbb{R}^{m \times n}$  is a diagonal rectangular matrix such that  $\Sigma_{i,i} \neq 0$  if and only if  $i \leq \text{rank}(\mathbf{A})$

The form  $\mathbf{A} = \mathbf{U}\mathbf{\Sigma}\mathbf{V}^T$  is called the Singular-Value Decomposition (SVD) of  $\mathbf{A}$ .

**Property 2.** Let  $r$  be the rank of a matrix  $\mathbf{A} \in \mathbb{R}^{m \times n}$ . Let  $\mathbf{U}$  and  $\mathbf{V}$  be the matrices from the SVD of  $\mathbf{A}$  with columns  $\mathbf{u}_1, \dots, \mathbf{u}_m \in \mathbb{R}^m$  and  $\mathbf{v}_1, \dots, \mathbf{v}_n \in \mathbb{R}^n$  respectively. Then the following hold:

- $\mathbf{u}_1, \dots, \mathbf{u}_r$  forms a basis a basis of  $\mathcal{R}(\mathbf{A})$
- $\mathbf{u}_{r+1}, \dots, \mathbf{u}_m$  forms a basis of  $\mathcal{N}(\mathbf{A}^T)$
- $\mathbf{v}_1, \dots, \mathbf{v}_r$  forms a basis a basis of  $\mathcal{R}(\mathbf{A}^T)$
- $\mathbf{v}_{r+1}, \dots, \mathbf{v}_n$  forms a basis of  $\mathcal{N}(\mathbf{A})$

**Definition 3 (Truncated SVD).** Theorem 1 can be rewritten as:

$$\mathbf{A} = \begin{matrix} \mathbf{U} & \mathbf{\Sigma} & \mathbf{V}^T \\ (m \times n) & (m \times m)(m \times n)(n \times n) & \end{matrix} = \begin{bmatrix} \tilde{\mathbf{U}} & \tilde{\mathbf{U}}_{\perp} \\ (m \times r) & (m \times m-r) \end{bmatrix} \begin{bmatrix} \mathbf{\Lambda} & \mathbf{0} \\ (r \times r) & (r \times n-r) \\ \mathbf{0} & \mathbf{0} \\ (m-r \times r) & (m-r \times n-r) \end{bmatrix} \begin{bmatrix} \tilde{\mathbf{V}}^T \\ (r \times n) \\ \tilde{\mathbf{V}}_{\perp}^T \\ (n-r \times n) \end{bmatrix} = \begin{matrix} \tilde{\mathbf{U}} & \mathbf{\Lambda} & \tilde{\mathbf{V}}^T \\ (m \times r) & (r \times r) & (r \times n) \end{matrix}$$

Where:

- $\mathbf{\Lambda} \in \mathbb{R}^{r \times r}$  is diagonal with  $\Lambda_{i,i} \neq 0$
- $\tilde{\mathbf{U}}^T\tilde{\mathbf{U}} = \tilde{\mathbf{V}}^T\tilde{\mathbf{V}} = \mathbf{I}_r$  ( $\tilde{\mathbf{U}}$  and  $\tilde{\mathbf{V}}$  are both orthogonal matrices)

The form  $\mathbf{A} = \tilde{\mathbf{U}}\mathbf{\Lambda}\tilde{\mathbf{V}}^T$  is called the Truncated Singular-Value Decomposition (Truncated SVD) of  $\mathbf{A}$ .

Remark : In truncated SVD,  $\tilde{\mathbf{U}}$  is such that  $\tilde{\mathbf{U}}^T \tilde{\mathbf{U}} = \mathbf{I}_r$ . However, it is not true in general that  $\tilde{\mathbf{U}} \tilde{\mathbf{U}}^T = \mathbf{I}_m$ . If  $r < m$ , we actually know that  $\tilde{\mathbf{U}} \tilde{\mathbf{U}}^T \neq \mathbf{I}_m$  because  $\mathbf{I}_m$  is of rank  $m$ , and cannot be linearly generated by  $\tilde{\mathbf{U}}$  of rank  $r < m$ .

$\mathbf{U}$  is orthogonal, so we have:

$$\mathbf{U}^T \mathbf{U} = \begin{bmatrix} \tilde{\mathbf{U}}^T \\ \tilde{\mathbf{U}}_\perp^T \end{bmatrix} \begin{bmatrix} \tilde{\mathbf{U}} & \tilde{\mathbf{U}}_\perp \end{bmatrix} = \begin{bmatrix} \tilde{\mathbf{U}}^T \tilde{\mathbf{U}} & \tilde{\mathbf{U}}^T \tilde{\mathbf{U}}_\perp \\ \tilde{\mathbf{U}}_\perp^T \tilde{\mathbf{U}} & \tilde{\mathbf{U}}_\perp^T \tilde{\mathbf{U}}_\perp \end{bmatrix} = \begin{bmatrix} \mathbf{I}_r & \mathbf{0} \\ \mathbf{0} & \mathbf{I}_{m-r} \end{bmatrix}$$

The same holds for  $\mathbf{V}^T \mathbf{V}$  with  $n$  instead of  $m$ .

### 3 Orthogonality and Projections

**Definition 4** (The dot product). *The dot product on  $\mathbb{R}^n$  has the following properties:*

- The dot product  $\langle \mathbf{a}, \mathbf{b} \rangle := \mathbf{a}^T \mathbf{b}$  is an inner product on  $\mathbb{R}^n$ .
- Two vectors  $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$  are orthogonal iff  $\langle \mathbf{a}, \mathbf{b} \rangle = 0$ .
- The inner product induces a norm on  $\mathbb{R}^n$ :  $\|\mathbf{v}\| = \sqrt{\langle \mathbf{v}, \mathbf{v} \rangle}$ .
- A basis  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k \in \mathbb{R}^n$  of a sub-space  $\mathcal{U}$  is orthonormal iff  $\begin{cases} \langle u_i, u_j \rangle = 0 & \forall i \neq j \\ \sqrt{\langle u_i, u_i \rangle} = \|u_i\| = 1 & \forall i \end{cases}$

**Definition 5** (Orthogonal matrix). *A matrix  $\mathbf{U} \in \mathbb{R}^{n \times k}$  is orthogonal if its columns form an orthonormal basis of  $\mathcal{R}(\mathbf{U})$ . Equivalently  $\mathbf{U}$  is orthogonal if and only if  $\mathbf{U}^T \mathbf{U} = \mathbf{I}$ .*

Note:  $\mathbf{U} \mathbf{U}^T$  is not necessarily equal to the identity matrix:  $\mathbf{U} \mathbf{U}^T \neq \mathbf{I}$ . However, if the matrix  $\mathbf{U}$  is square i.e.  $\mathbf{U} \in \mathbb{R}^{n \times n}$ , then  $\mathbf{U}^T \mathbf{U} = \mathbf{I}$  implies  $\mathbf{U} \mathbf{U}^T = \mathbf{I}$

Remark: if  $\mathbf{U} \in \mathbb{R}^{n \times k}$  is orthogonal, then  $\|\mathbf{U} \mathbf{x}\| = \|\mathbf{x}\|$ . This is easily seen that this follows from the orthogonality of the matrix  $\mathbf{U}$ :  $\|\mathbf{U} \mathbf{x}\|^2 = \mathbf{x}^T \mathbf{U}^T \mathbf{U} \mathbf{x} = \mathbf{x}^T \mathbf{x} = \|\mathbf{x}\|^2$ .

**Property 6.** *If  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k \in \mathbb{R}^n$  is an orthonormal basis of a sub-space  $\mathcal{U}$ ,*

$$\mathbf{U} = \begin{bmatrix} | & | & \dots & | \\ \mathbf{u}_1 & \mathbf{u}_2 & \dots & \mathbf{u}_k \\ | & | & \dots & | \end{bmatrix} \in \mathbb{R}^{n \times k} \Rightarrow \forall \mathbf{x} \in \mathcal{U}, \mathbf{x} = \mathbf{U} \mathbf{U}^T \mathbf{x}$$

Consequently  $\|\mathbf{x}\| = \|\mathbf{U}^T \mathbf{x}\|$ .

Note: Even though  $\mathbf{U} \mathbf{U}^T$  is not the identity matrix, it acts as such on the sub-space  $\mathcal{U}$ .

*Proof.* A vector  $\mathbf{x}$  in the sub-space  $\mathcal{U}$  can, by definition, be written as a linear combination of its basis vectors.


$$\mathbf{x} \in \mathcal{U} \Rightarrow \exists \mathbf{a} \in \mathbb{R}^k : \mathbf{x} = \mathbf{U} \mathbf{a}$$

To prove that  $\mathbf{U} \mathbf{U}^T$  acts as an identity matrix, we look at what it does to  $\mathbf{x}$ . Using the fact that  $\mathbf{x}$  can be written as a linear combination of the basis vectors of the sub-space  $\mathcal{U}$ , and that  $\mathbf{U}$  is defined as an orthogonal matrix, we have:

$$\mathbf{U} \mathbf{U}^T \mathbf{x} = \mathbf{U} \mathbf{U}^T \mathbf{U} \mathbf{a} = \mathbf{U} \mathbf{a} = \mathbf{x}.$$

A consequence of this property is that  $\mathbf{x}$  and  $\mathbf{U}^T \mathbf{x}$  are vectors of the same size:

$$\|\mathbf{x}\|^2 = \mathbf{x}^T \mathbf{x} = (\mathbf{U} \mathbf{U}^T \mathbf{x})^T \mathbf{x} = \mathbf{x}^T \mathbf{U} \mathbf{U}^T \mathbf{x} = (\mathbf{U}^T \mathbf{x})^T (\mathbf{U}^T \mathbf{x}) = \|\mathbf{U}^T \mathbf{x}\|^2 \quad \square$$

Figure 1: Orthogonal projection of a vector  $\mathbf{x}$  onto a sub-space  $\mathcal{U}$ 

**Definition 7** (Orthogonal Projection). Let  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k \in \mathbb{R}^n$  be an orthonormal basis of some sub-space  $\mathcal{U}$  and

$$U = \begin{bmatrix} | & | & \cdots & | \\ \mathbf{u}_1 & \mathbf{u}_2 & \cdots & \mathbf{u}_k \\ | & | & \cdots & | \end{bmatrix} \in \mathbb{R}^{n \times k}$$

The orthogonal projection onto  $\mathcal{U}$  is defined as:

$$\begin{aligned} \Pi_{\mathcal{U}} : \mathbb{R}^n &\rightarrow \mathbb{R}^n \\ \mathbf{x} &\rightarrow \mathbf{U}\mathbf{U}^T\mathbf{x} \end{aligned}$$

This is well defined and unique. In other words, if there is another orthogonal matrix  $\mathbf{V} \in \mathbb{R}^{n \times k}$  such that  $\mathcal{R}(\mathbf{V}) = \mathcal{U}$ , then we can prove that the two projections are the same:  $\mathbf{U}\mathbf{U}^T = \mathbf{V}\mathbf{V}^T$ .

*Proof.* Since the columns of  $\mathbf{U}$  and  $\mathbf{V}$  consist of the orthonormal basis vectors of the sub-space  $\mathcal{U}$ , they span the same sub-space i.e. they have the same range:  $\mathcal{R}(\mathbf{V}) = \mathcal{R}(\mathbf{U})$ .

But if this is true, then there must exist a linear map  $\mathbf{P}$  between the two matrices:

$$\exists \mathbf{P} \in \mathbb{R}^{k \times k} \text{ such that } \mathbf{V} = \mathbf{U}\mathbf{P}$$

But since the matrices  $\mathbf{U}$  and  $\mathbf{V}$  are orthogonal, we have:

$$\mathbf{I} = \mathbf{V}^T\mathbf{V} = \mathbf{P}^T\mathbf{U}^T\mathbf{U}\mathbf{P} = \mathbf{P}^T\mathbf{P}$$

So we have proven that  $\mathbf{P}$  is also orthogonal. Since  $\mathbf{P}$  is a square matrix, we have that:  $\mathbf{P}^T\mathbf{P} = \mathbf{P}\mathbf{P}^T = \mathbf{I}$

Inserting this identity matrix below, we prove that orthogonal projection is unique.

$$\mathbf{V}\mathbf{V}^T = \mathbf{U}\mathbf{P}\mathbf{P}^T\mathbf{U}^T = \mathbf{U}\mathbf{U}^T \quad \square$$

**Definition 8** (Orthogonal Complement). If  $\mathcal{U}$  is a sub-space of  $\mathbb{R}^n$ , the orthogonal complement of  $\mathcal{U}$  is defined as:  $\mathcal{U}^\perp = \{\mathbf{v} \in \mathbb{R}^n \mid \langle \mathbf{u}, \mathbf{v} \rangle = 0 \forall \mathbf{u} \in \mathcal{U}\}$ .


**Property 9.**  $\mathcal{U}^\perp$  is a sub-space of  $\mathbb{R}^n$  and  $\mathbb{R}^n = \mathcal{U} \oplus \mathcal{U}^\perp$ .

*Proof.* Let  $\mathbf{u}, \mathbf{v} \in \mathcal{U}^\perp$ . For any  $\mathbf{w} \in \mathcal{U}$  we have  $\langle \mathbf{w}, \mathbf{u} + \mathbf{v} \rangle = \langle \mathbf{w}, \mathbf{u} \rangle + \langle \mathbf{w}, \mathbf{v} \rangle = 0 + 0 = 0$ , so  $\mathbf{u} + \mathbf{v} \in \mathcal{U}^\perp$ .

Let  $\mathbf{u} \in \mathcal{U}^\perp$  and  $\alpha \in \mathbb{R}$ . Then for any  $\mathbf{w} \in \mathcal{U}$  we have  $\langle \mathbf{w}, \alpha\mathbf{u} \rangle = \alpha\langle \mathbf{w}, \mathbf{u} \rangle = \alpha \cdot 0 = 0$ , so  $\alpha\mathbf{u} \in \mathcal{U}^\perp$ .

Therefore  $\mathcal{U}^\perp$  is a sub-space.

We also know that zero must be a part of every sub-space. In fact, zero is the only common element of these two sub-spaces (Assuming that there is a non-zero vector  $\mathbf{w}$  that belongs to both of the sub-spaces  $\mathcal{U}$  and  $\mathcal{U}^\perp$ , using the above definition of  $\mathcal{U}^\perp$ , it is easily seen that the inner product of  $\mathbf{w}$  with itself is 0 i.e.  $\langle \mathbf{w}, \mathbf{w} \rangle = 0$  which is contradictory to the assumption that  $\mathbf{w} \neq 0$ .) A vector  $\mathbf{x} \in \mathbb{R}^n$  can be written as  $\mathbf{x} = \Pi_{\mathcal{U}}(\mathbf{x}) + \mathbf{v}$ . It can be proven (see property 10 below) that the vector  $\mathbf{v} = \mathbf{x} - \Pi_{\mathcal{U}}(\mathbf{x})$  must be orthogonal to  $\Pi_{\mathcal{U}}(\mathbf{x}) \in \mathcal{U}$  and that it must, therefore, belong to the orthogonal complement of  $\mathcal{U}$ . So  $\mathbb{R}^n = \mathcal{U} + \mathcal{U}^\perp$ .

Figure 2: Splitting a vector into its components on two sub-spaces  $\mathcal{U}$  and  $\mathcal{U}^\perp$ 

Since 0 is the only vector in common of  $\mathcal{U}$  and  $\mathcal{U}_\perp$ ,  $\mathbb{R}^n = \mathcal{U} \oplus \mathcal{U}^\perp$ , i.e. every vector in  $\mathbb{R}^n$  can be decomposed uniquely into a sum of two vectors, one belonging to  $\mathcal{U}$  and the other to  $\mathcal{U}^\perp$ .  $\square$

**Property 10.** Based on the above definitions of orthogonality and orthogonal complement, we can summarize the following properties of orthogonal projection:

- $\Pi_{\mathcal{U}}^2 = \Pi_{\mathcal{U}}$
- $\forall \mathbf{x} : \langle \Pi_{\mathcal{U}}(\mathbf{x}), \Pi_{\mathcal{U}}(\mathbf{x}) - \mathbf{x} \rangle = 0$
- $\text{Im}(\Pi_{\mathcal{U}}) = \mathcal{U}$
- $\text{Ker}(\Pi_{\mathcal{U}}) = \mathcal{U}^\perp$
- $\forall \mathbf{x} : \|\Pi_{\mathcal{U}}(\mathbf{x})\| \leq \|\mathbf{x}\|$

*Proof.* These properties can be proven as follows :

- $\Pi_{\mathcal{U}}^2 = \Pi_{\mathcal{U}}$ :

To prove this we use the definition of projection in terms of matrices: Knowing that  $\mathbf{U}\mathbf{U}^T\mathbf{x}$  is a projection, we have:  $\Pi_{\mathcal{U}}^2(\mathbf{x}) = \Pi_{\mathcal{U}}(\mathbf{U}\mathbf{U}^T\mathbf{x}) = \mathbf{U}\mathbf{U}^T\mathbf{U}\mathbf{U}^T\mathbf{x} = \mathbf{U}\mathbf{U}^T\mathbf{x} = \Pi_{\mathcal{U}}(\mathbf{x})$ .

Note that in this calculation we used  $\mathbf{U}^T\mathbf{U} = \mathbf{I}$ .

- $\forall \mathbf{x} : \langle \Pi_{\mathcal{U}}(\mathbf{x}), \Pi_{\mathcal{U}}(\mathbf{x}) - \mathbf{x} \rangle = 0$  :

$$\begin{aligned} \langle \Pi_{\mathcal{U}}(\mathbf{x}), \Pi_{\mathcal{U}}(\mathbf{x}) - \mathbf{x} \rangle &= \langle \mathbf{U}\mathbf{U}^T\mathbf{x}, \mathbf{U}\mathbf{U}^T\mathbf{x} - \mathbf{x} \rangle \\ &= \mathbf{x}^T\mathbf{U}\mathbf{U}^T\mathbf{U}\mathbf{U}^T\mathbf{x} - \mathbf{x}^T\mathbf{U}\mathbf{U}^T\mathbf{x} \\ &= 0 \end{aligned}$$

- $\text{Im}(\Pi_{\mathcal{U}}) = \mathcal{U}$ :

$\forall \mathbf{x} \in \mathbb{R}^n$  we have that  $\Pi_{\mathcal{U}}(\mathbf{x}) \in \mathcal{U}$  because since  $\mathbf{U}$  is a basis of  $\mathcal{U}$ ,  $\mathbf{U}\mathbf{U}^T\mathbf{x} \in \mathcal{U}$ , so  $\text{Im}(\Pi_{\mathcal{U}}) \subset \mathcal{U}$ .

If  $\mathbf{y} \in \mathcal{U}$  then  $\Pi_{\mathcal{U}}(\mathbf{y}) = \mathbf{U}\mathbf{U}^T\mathbf{y} = \mathbf{y}$  (see property 6), so  $\text{Im}(\Pi_{\mathcal{U}}) \supset \mathcal{U}$ .

So  $\text{Im}(\Pi_{\mathcal{U}}) = \mathcal{U}$

- $\text{Ker}(\Pi_{\mathcal{U}}) = \mathcal{U}_\perp$ :

Let  $\mathbf{x} \in \text{Ker}(\Pi_{\mathcal{U}})$ . Then  $\Pi_{\mathcal{U}}(\mathbf{x}) = 0$ , so  $\mathbf{U}\mathbf{U}^T\mathbf{x} = 0$ .

We know from property 9 that  $\mathbf{x}$  can be uniquely decomposed into  $\mathbf{x} = \mathbf{x}' + \mathbf{x}'_\perp$  where  $\mathbf{x}' \in \mathcal{U}$  and  $\mathbf{x}'_\perp \in \mathcal{U}_\perp$ .

So  $0 = \mathbf{U}\mathbf{U}^T\mathbf{x}' + \mathbf{U}\mathbf{U}^T\mathbf{x}'_\perp = \mathbf{x}' + \mathbf{U}\mathbf{U}^T\mathbf{x}'_\perp$ .

Since  $\mathbf{x}'_\perp \in \mathcal{U}_\perp$ ,  $\mathbf{U}\mathbf{U}^T\mathbf{x}'_\perp = 0$ , so  $\mathbf{x}' = 0$ .

Therefore  $\mathbf{x} = \mathbf{x}'_{\perp} \in \mathcal{U}_{\perp}$ , so  $\text{Ker}(\Pi_{\mathcal{U}}) \subset \mathcal{U}_{\perp}$ .

Let  $\mathbf{x} \in \mathcal{U}_{\perp}$ , then  $\forall \mathbf{u} \in \mathcal{U}, \langle \mathbf{u}, \mathbf{x} \rangle = 0$ , so  $\mathbf{U}\mathbf{U}^T \mathbf{x} = \mathbf{0}$ . So  $\text{Ker}(\Pi_{\mathcal{U}}) \supset \mathcal{U}_{\perp}$ .

So  $\text{Ker}(\Pi_{\mathcal{U}}) = \mathcal{U}_{\perp}$ .

- $\forall \mathbf{x} : \|\Pi_{\mathcal{U}}(\mathbf{x})\| \leq \|\mathbf{x}\|$ :

We know that any vector  $\mathbf{x}$  can be uniquely decomposed into two components: one is  $\Pi_{\mathcal{U}}(\mathbf{x}) \in \mathcal{U}$ , and the other is in  $\mathcal{U}_{\perp}$ . For simplicity we call the other component  $\mathbf{v}$ .

$$\mathbf{x} = \Pi_{\mathcal{U}}(\mathbf{x}) + \mathbf{v}$$

$$\|\mathbf{x}\|^2 = \|\Pi_{\mathcal{U}}(\mathbf{x}) + \mathbf{v}\|^2 = \langle \Pi_{\mathcal{U}}(\mathbf{x}) + \mathbf{v}, \Pi_{\mathcal{U}}(\mathbf{x}) + \mathbf{v} \rangle = \langle \Pi_{\mathcal{U}}(\mathbf{x}), \Pi_{\mathcal{U}}(\mathbf{x}) \rangle + \langle \Pi_{\mathcal{U}}(\mathbf{x}), \mathbf{v} \rangle + \langle \mathbf{v}, \Pi_{\mathcal{U}}(\mathbf{x}) \rangle + \langle \mathbf{v}, \mathbf{v} \rangle$$

Since the vector  $\mathbf{v}$  is in the orthogonal complement of  $\mathcal{U}$ , it must be orthogonal to all the vectors on  $\mathcal{U}$ .

$$\|\mathbf{x}\|^2 = \|\Pi_{\mathcal{U}}(\mathbf{x})\|^2 + \|\mathbf{v}\|^2 \Rightarrow \|\Pi_{\mathcal{U}}(\mathbf{x})\|^2 \leq \|\mathbf{x}\|^2 \Rightarrow \|\Pi_{\mathcal{U}}(\mathbf{x})\| \leq \|\mathbf{x}\|$$

□

**Property 11.** Let  $\mathcal{U}$  be a subspace of  $\mathbb{R}^n$ . Then, for any  $\mathbf{x} \in \mathbb{R}^n$  we have  $\arg \min_{\mathbf{u} \in \mathcal{U}} \|\mathbf{u} - \mathbf{x}\| = \Pi_{\mathcal{U}}(\mathbf{x})$ . I.e. the closest point to  $\mathbf{x}$  which belongs to  $\mathcal{U}$  is the orthogonal projection of  $\mathbf{x}$  onto  $\mathcal{U}$ .